

Exercises CFT-course fall 2011, set 7.

Due on december 2nd, 2011.

1. In this exercise, we will derive an ordinary differential equation for the four point function of $\phi_{1,2}$ of the minimal models:

$$G(\{w_i\}) = \langle \phi_{2,1}(w_1)\phi_{2,1}(w_2)\phi_{2,1}(w_3)\phi_{2,1}(w_4) \rangle = (w_1 - w_2)^{-2h}(w_3 - w_4)^{-2h} F(x) ,$$

where $x = \frac{(w_1-w_2)(w_3-w_4)}{(w_1-w_4)(w_3-w_2)}$, and h is the conformal dimension of $\phi_{2,1}$.

The level-2 null-vector condition translates into the following partial differential equations:

$$\left(\frac{3}{2(2h+1)} \partial_{w_i}^2 - \sum_{j \neq i} \left(\frac{\partial_{w_j}}{(w_i - w_j)} + \frac{h}{(w_i - w_j)^2} \right) \right) G(\{w_i\}) = 0 .$$

- a. Show, by considering the special values $z_1 = 0, z_2 = x, z_3 = \infty, z_4 = 1$, that $F(x)$ satisfies

$$\left(x(1-x)\partial_x^2 + \left(\frac{2(1-4h)}{3} + \frac{4(h-1)}{3}x \right) \partial_x - \frac{2h(2h+1)}{3} \frac{x}{1-x} \right) F(x) = 0$$

- b. Show that the differential equation obtained in a. can be brought into the form of the hypergeometric differential equation, by defining a function $H(x) = f(x)F(x)$, such that $H(x)$ satisfies

$$\left(x(1-x)\partial_x^2 + \left(\frac{2(1-4h)}{3} + \frac{4(4h-1)}{3}x \right) \partial_x + \frac{4h(1-4h)}{3} \right) H(x) = 0 \quad (1)$$

2. Show that in the case of the Ising model, two independent solutions of the differential equation (1) (with $h = \frac{1}{16}$) take the form

$$H^\pm(x) = \sqrt{\frac{1 \pm \sqrt{1-x}}{2}} ,$$

by making a suitable (trigonometric) transformation.

Thus, the (chiral) four-point σ -correlator of the Ising model takes the form

$$\langle \sigma(w_1)\sigma(w_2)\sigma(w_3)\sigma(w_4) \rangle^\pm = (w_1 - w_2)^{-1/8}(w_3 - w_4)^{-1/8}(1-x)^{-1/8} \sqrt{\frac{1 \pm \sqrt{1-x}}{2}} .$$

3. Majorana fermions with periodic and anti-periodic boundary conditions.

The mode expansion for free (Majorana) fermions reads $\psi(z) = \sum_n \psi_n z^{-n-1/2}$, or $\psi_n = \oint \frac{dz}{2\pi i} z^{n-1/2} \psi(z)$.

- a. Show that the modes obey $\{\psi_n, \psi_m\} = \delta_{n+m,0}$.

We will now consider periodic and anti-periodic boundary conditions for the fermion $\psi(z)$ when z is moved around the origin: $\psi(e^{2\pi i}z) = \pm\psi(z)$. The modes n are half integer $n \in \mathbb{Z} + \frac{1}{2}$ in the periodic (P) case, and integer $n \in \mathbb{Z}$ in the anti-periodic (A) case.

- b. Use the explicit mode expansions to show that

$$\langle\psi(z)\psi(w)\rangle_{\text{P}} = \frac{1}{z-w}$$

$$\langle\psi(z)\psi(w)\rangle_{\text{A}} = \frac{1}{2} \frac{\sqrt{\frac{z}{w}} + \sqrt{\frac{w}{z}}}{z-w}$$

It is given that

$$\langle\sigma(w_1)\psi(z_1)\psi(z_2)\sigma(w_2)\rangle = \frac{1}{2}(w_1 - w_2)^{-1/8} \frac{\left(\frac{(z_1-w_1)(z_2-w_2)}{(z_1-w_2)(z_2-w_1)}\right)^{1/2} + \left(\frac{(z_1-w_2)(z_2-w_1)}{(z_1-w_1)(z_2-w_2)}\right)^{1/2}}{(z_1 - z_2)}$$

- c. Consider $\langle\sigma(\infty)|\psi(z_1)\psi(z_2)|\sigma(0)\rangle$, where $\langle\sigma(\infty)| = \lim_{w \rightarrow \infty} \langle 0|\sigma(w)w^{2h}$, and argue that σ ‘changes the boundary conditions on ψ ’.