

Exercises CFT-course fall 2011, set 9.

Due on friday, december 16th.

1. CFT description of the transverse-field Ising model.

In exercise 1. of the first set of exercises, we considered the transverse-field Ising model in one dimension. The hamiltonian in terms of the Pauli matrices σ was given by

$$H_I = -J \sum_i (g\sigma_i^x + \sigma_i^z \sigma_{i+1}^z) .$$

You showed that the hamiltonian can be diagonalized, with as the final result

$$H_I = \sum_k \varepsilon_k \left(\gamma_k^\dagger \gamma_k - \frac{1}{2} \right) \quad \varepsilon_k = 2(1 + g^2 - 2g \cos(ak))^{1/2}$$

Here, the $\gamma_k^{(\dagger)}$'s are fermion creation and annihilation operators with momentum k . For F odd, the momenta take the values $k = 0, 1, \dots, L-1$, while for F even, the allowed momenta are $k = 1/2, 3/2, \dots, L-1/2$, because of the periodic boundary conditions we imposed.

In this exercise, we will consider the critical behaviour at $g = 1$, for $J = 1$. Although the model has been solved exactly, it is not so easy to extract the critical behaviour analytically, because of the sum over k , which one has to evaluate to obtain the energies. Therefore, the use of some computer program to evaluate the energies for finite system size.

- Evaluate the the ground state energy (at $g = J = 1$) numerically for finite system sizes L , (up to $L = 500$ should not be a problem) and fit the results to the (general) finite size scaling formula

$$E_i = E_0 L + \frac{2\pi v}{L} \left(-\frac{c}{12} + h_l + h_r \right) ,$$

where E_0 is a constant energy per spin, v a velocity, c the central charge, and h_l and h_r the left and right scaling dimensions of scaling fields.

- Show numerically that the system is critical, by obtaining the finite size gap between the ground and first excited states.
- Obtain the other constants in the scaling formula and dimensions of the primary fields by considering various low-lying states in the spectrum.
- (optional) Obtain the full spectrum for a small system (say $L = 10$), and plot the states as a function of momentum.

(PTO)

2. Entanglement entropy from conformal field theory.

Study the paper ‘Entanglement entropy and conformal field theory’ by P. Calabrese and J. Cardy (J.Phys. A**42**, 504005 (2009); arXiv:0905.4013) up to (at least) section 3.2.

Briefly describe the structure of the calculation, and fill in the details which were skipped during the lecture, in particular the calculation of $\langle T(w) \rangle_{\mathcal{R}_n}$ and the correlator $\langle \Phi_n(u, 0) \Phi_{-n}(v, 0) \rangle_{\mathcal{L}^{(n)}, \mathbf{C}}$.