

## Exercises CFT-course fall 2013, set 8.

Due on december 6th, 2013.

1. In this exercise, we will derive an ordinary differential equation for the four point function of  $\phi_{2,1}$  of the minimal models:

$$G(\{w_i\}) = \langle \phi_{2,1}(w_1)\phi_{2,1}(w_2)\phi_{2,1}(w_3)\phi_{2,1}(w_4) \rangle = (w_1 - w_2)^{-2h}(w_3 - w_4)^{-2h}F(x) ,$$

where  $x = \frac{(w_1-w_2)(w_3-w_4)}{(w_1-w_4)(w_3-w_2)}$ , and  $h$  is the conformal dimension of  $\phi_{2,1}$ .

The level-2 null-vector condition translates into the following partial differential equations:

$$\left( \frac{3}{2(2h+1)}\partial_{w_i}^2 - \sum_{j \neq i} \left( \frac{\partial_{w_j}}{(w_i - w_j)} + \frac{h}{(w_i - w_j)^2} \right) \right) G(\{w_i\}) = 0 .$$

- a. Show, by considering the special values  $w_1 = 0, w_2 = x, w_3 = \infty, w_4 = 1$ , that  $F(x)$  satisfies

$$\left( x(1-x)\partial_x^2 + \left( \frac{2(1-4h)}{3} + \frac{4(h-1)}{3}x \right) \partial_x - \frac{2h(2h+1)}{3} \frac{x}{1-x} \right) F(x) = 0$$

- b. Show that the differential equation obtained in a. can be brought into the form of the hypergeometric differential equation, by defining a function  $H(x) = f(x)F(x)$ , such that  $H(x)$  satisfies

$$\left( x(1-x)\partial_x^2 + \left( \frac{2(1-4h)}{3} + \frac{4(4h-1)}{3}x \right) \partial_x + \frac{4h(1-4h)}{3} \right) H(x) = 0 \quad (1)$$

2. Show that in the case of the Ising model, two independent solutions of the differential equation (1) (with  $h = \frac{1}{16}$ ) take the form

$$H^\pm(x) = \sqrt{\frac{1 \pm \sqrt{1-x}}{2}} ,$$

by making a suitable (trigonometric) transformation.

Thus, the (chiral) four-point  $\sigma$ -correlator of the Ising model takes the form

$$\langle \sigma(w_1)\sigma(w_2)\sigma(w_3)\sigma(w_4) \rangle^\pm = (w_1 - w_2)^{-1/8}(w_3 - w_4)^{-1/8}(1-x)^{-1/8} \sqrt{\frac{1 \pm \sqrt{1-x}}{2}} .$$

3. Majorana fermions with periodic and anti-periodic boundary conditions.

The mode expansion for free (Majorana) fermions reads  $\psi(z) = \sum_n \psi_n z^{-n-1/2}$ , or  $\psi_n = \oint \frac{dz}{2\pi i} z^{n-1/2} \psi(z)$ .

- a. Show that the modes obey  $\{\psi_n, \psi_m\} = \delta_{n+m,0}$ .

We will now consider periodic and anti-periodic boundary conditions for the fermion  $\psi(z)$  when  $z$  is moved around the origin:  $\psi(e^{2\pi i} z) = \pm\psi(z)$ . The modes  $n$  are half integer  $n \in \mathbb{Z} + \frac{1}{2}$  in the periodic (P) case, and integer  $n \in \mathbb{Z}$  in the anti-periodic (A) case.

- b. Use the explicit mode expansions to show that

$$\langle \psi(z)\psi(w) \rangle_{\text{P}} = \frac{1}{z-w}$$

$$\langle \psi(z)\psi(w) \rangle_{\text{A}} = \frac{1}{2} \frac{\sqrt{\frac{z}{w}} + \sqrt{\frac{w}{z}}}{z-w}$$

It is given that

$$\langle \sigma(w_1)\psi(z_1)\psi(z_2)\sigma(w_2) \rangle = \frac{1}{2}(w_1 - w_2)^{-1/8} \frac{\left(\frac{(z_1-w_1)(z_2-w_2)}{(z_1-w_2)(z_2-w_1)}\right)^{1/2} + \left(\frac{(z_1-w_2)(z_2-w_1)}{(z_1-w_1)(z_2-w_2)}\right)^{1/2}}{(z_1 - z_2)}$$

- c. Consider  $\langle \sigma(\infty)|\psi(z_1)\psi(z_2)|\sigma(0) \rangle$ , where  $\langle \sigma(\infty)| = \lim_{w \rightarrow \infty} \langle 0|\sigma(w)w^{2h}$ , and argue that  $\sigma$  ‘changes the boundary conditions on  $\psi$ ’.