

## Exercises CFT-course fall 2018, set 4.

Due on friday, november 30, 2018.

1. The two point function on the cylinder.
  - a. Calculate the two point function of a primary operator  $\phi(z, \bar{z})$  with scaling dimensions  $(h, \bar{h})$  on the cylinder, by using the conformal map  $w = \frac{L \log z}{2\pi}$ . Result:

$$\langle \phi(w_1, \bar{w}_1) \phi(w_2, \bar{w}_2) \rangle = \left( \frac{2\pi}{L} \right)^{2h+2\bar{h}} \left( 2 \sinh(\pi(w_1 - w_2)/L) \right)^{-2h} \left( 2 \sinh(\pi(\bar{w}_1 - \bar{w}_2)/L) \right)^{-2\bar{h}}$$

- b. Let  $w = t + iu$ . Analyze the result of a. in the limits  $t_1 - t_2 \ll L$  and  $t_1 - t_2 \gg L$  and discuss the result. For simplicity, you may assume that  $h = \bar{h}$ .
2. The massless free boson  $\phi$  has the action

$$S = \frac{g}{2} \int dx^2 \partial_\mu \phi \partial^\mu \phi .$$

- a. Determine the propagator of the free massless boson theory:

$$\langle \phi(z, \bar{z}) \phi(w, \bar{w}) \rangle = \frac{-1}{4\pi g} (\ln(z - w) + \ln(\bar{z} - \bar{w})) + \text{const.}$$

- b. Determine the form of the stress-energy tensor, namely  $T(z) = -2\pi g : \partial\phi(z)\partial\phi(z) :$ .
  - c. Calculate the OPE  $T(z)T(w)$ .
  - d. Show that the vertex operator  $V = : e^{i\alpha\phi} : (z)$  is a primary field of dimension  $h_\alpha = \frac{\alpha^2}{8\pi g}$ .

3. The massless free fermion.

The action of the massless free fermion can be written as

$$S = g \int dx^2 (\bar{\psi} \partial \bar{\psi} + \psi \bar{\partial} \psi) ,$$

where  $\psi$  and  $\bar{\psi}$  are the two components of the spinor  $\Psi$ . It is given that the equations of motion are  $\partial \bar{\psi} = \bar{\partial} \psi = 0$ , while the propagator takes the form  $\langle \psi(z, \bar{z}) \psi(w, \bar{w}) \rangle = \frac{1}{2\pi g} \frac{1}{z-w}$ . The stress-energy tensor can be written as  $T(z) = -\pi g : \psi(z) \partial \psi(z) :$

- a. Show that  $\psi(z)$  is a primary field.
  - b. Calculate the OPE  $T(z)T(w)$ .