

# A little on chaos, thermalisation, localisation... (well, dynamics in general) in quantum systems

**Jonas Larson**

Stockholm University and Universität zu Köln

York 21/7-2015



# Motivation

- Long time evolution of closed quantum systems not fully understood.
- Cold atom system → Not only of academic interest.
- Open questions in closed system quantum dynamics:
  - i. Criteria for equilibration/thermalization.
  - ii. Mechanism behind thermalization.
  - iii. Properties of equilibrated states.
  - iv. Definition for “quantum integrability”.
  - v. Many-body localization...
  - vi. Open systems...

# Before take-off

- Interest in many-body systems: Large hilbert spaces - difficult to handle on a classical computer. 1 mole → hilbert space dimension  $2^{10^{23}}$ , number of atoms in the universe  $10^{80}$ .
- Often universal (underlying symmetries) properties of hamiltonians (Random matrix theory) - independent of number of degrees of freedom.
- Large matrices also with few degrees of freedom, current case.

# Outline

1. Quantum Thermalization.
2. Integrability. Problems.
3. Chaos. Problems.
4. Localization and absence of *ETH*.



# Quantum Thermalization



# Quantum Thermalization

- Characteristics of  $\hat{\rho}$  for long times.

- 1) Weak coupling.
- 2) Infinite degrees of freedom (bath).
- 3) Delta correlated in time (bath): Markov approximation (no memory).
- 4) Factorizable system-bath state (Born approximation).

Open system

Thermalization of system.




$$\partial_t \hat{\rho} = i[\hat{\rho}, \hat{H}_{sys}] + \hat{L}[\hat{\rho}]$$

# Quantum Thermalization

## Equilibration

Closed system

- Characteristics of  $\hat{\rho}_S(t)$  for long times.
- No clear separation “system/bath”, no Born-Markov nor rotating-wave approximations.



# Quantum Thermalization

## Equilibration

Closed system

- Characteristics of  $\hat{\rho}(t)$  for long times.
- Equilibration:

$$\langle \hat{A} \rangle = \text{Tr}[\hat{A}\hat{\rho}(t)], \quad \begin{cases} t - \text{independent as } t \rightarrow \infty \\ \hat{A} \text{ local observable.} \end{cases}$$

- Thermalization:

$$\langle \hat{A} \rangle = \langle \hat{A} \rangle_{Th} \quad \text{at long times}$$

$$\langle \hat{A} \rangle_{Th} = \text{Tr}[\hat{A}\hat{\rho}_{Th}],$$

where  $\hat{\rho}_{Th}$  = thermal state. “Temperature” determined from  $\langle \hat{H} \rangle$ .  
**No** memory of initial state.



# Quantum Thermalization

## ETH – *Eigenstate Thermalization Hypotesis*

- $|\Psi(t)\rangle = \sum_{\gamma} C_{\gamma} e^{-iE_{\gamma}t/\hbar} |\psi_{\gamma}\rangle \rightarrow$

$$\langle \hat{A}(t) \rangle = \sum_{\gamma, \delta} C_{\delta}^* C_{\gamma} e^{i(E_{\delta} - E_{\gamma})t/\hbar} A_{\delta\gamma}, \quad A_{\delta\gamma} = \langle \psi_{\delta} | \hat{A} | \psi_{\gamma} \rangle$$

- If thermalization (long-time limit)

$$\langle \hat{A} \rangle^{LT} = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T dt \langle \hat{A}(t) \rangle = \sum_{\gamma} |C_{\gamma}|^2 A_{\gamma\gamma}.$$

- *ETH*:  $A_{\gamma\gamma}$  is approximately constant in the "energy window" of the state  $\Psi$ .
- *ETH*: For all  $\gamma$ ,  $\hat{\rho}_A = \text{Tr}_B [|\psi_{\gamma}\rangle\langle\psi_{\gamma}|]$  is thermal.

# Quantum Thermalization

## Thermalization

- Which systems thermalize?

Possible candidates:

- 1) Quantum non-integrable systems.
- 2) Chaotic systems.



# Integrability

# Integrability

## 101 Quantum Integrability

- Classical systems:

**Definition:** A system is integrable if the number of degrees of freedom  $N$  is smaller than or equal to the number  $K$  of *independent* constants of motion.

$$\{Q_n, H\} = 0, \quad n = 1, 2, \dots, K, \quad \{Q_n, Q_m\} = 0 \quad \forall n, m$$

# Integrability

## 101 Quantum Integrability

- Quantum systems:

**Definition 1:** Replace  $\{ , \} \rightarrow i[ , ]/\hbar$ . Fails, take  $\hat{P}_\gamma = |\Psi_\gamma\rangle\langle\Psi_\gamma|$ .

**Definition 2:** Use definition 1, but consider *relevant* constants of motion - that is operators with classical counterparts. Fails, not all operators have any classical corresponding observable.

**Definition 3:** Poissonian level statistics ( $P(S) = e^{-S}$ ) implies integrability.

**Definition 4:** Level crossings implies integrability.

⋮

**Definition 64:** A quantum system is integrable if it is exactly solvable.

# Integrability vs thermalization

## Spin-orbit coupled particle

Rabi Model

- *Rabi Hamiltonian* of quantum optics

$$\hat{H}_R = \omega \hat{a}^\dagger \hat{a} + \frac{\Omega}{2} \hat{\sigma}_z + v(\hat{a}^\dagger + \hat{a}) \hat{\sigma}_x.$$

- $Z_2$ -parity symmetry

$$[\hat{U}_p, \hat{H}_R] = 0, \quad \hat{U}_p = e^{i\pi(\hat{a}^\dagger \hat{a} + \frac{\hat{\sigma}_z}{2})}.$$

- Drive term breaks  $Z_2$  (total energy only preserved quantity)


$$\hat{H}_{dR} = \omega \hat{a}^\dagger \hat{a} + \frac{\Omega}{2} \hat{\sigma}_z + v(\hat{a}^\dagger + \hat{a}) \hat{\sigma}_x + \gamma \hat{\sigma}_x.$$

# Integrability vs thermalization

## Spin-orbit coupled particle

Rabi Model

- Is the driven Rabi model integrable?
- **Definition 3:** Level-statistics. Two branches, neither Poissonian  
→ Non-integrable.




Level statistics of the Rabi model.

# Integrability vs thermalization

## Spin-orbit coupled particle

Rabi Model

- Is the driven Rabi model integrable?
- **Definition 4:** Avoided crossings. No visible crossings  $\rightarrow$  Non-integrable.



Energies of the Rabi model.



# Integrability vs thermalization

## Spin-orbit coupled particle

- Is the driven Rabi model integrable?
- **Definition 64**: Solvable. Braak (PRL 2011) says it might be solvable but not integrable, others say it is *quasi solvable* → integrable?


Rabi Model

# Integrability vs thermalization

## Spin-orbit coupled particle

Rabi Model

- Does the driven Rabi model thermalize?
- If quantum non-integrability implies thermalization a qualified guess would be yes.



Scaled variance of  $\langle \hat{n}(t) \rangle$ . Thermalization  $\rightarrow \delta_n = 0$ . No thermalization!



# Chaos

# Classical chaos

## Butterfly effect

*Hamilton equations:*

$$\frac{dp_j}{dt} = -\frac{\partial H}{\partial q_j}, \quad \frac{dq_j}{dt} = \frac{\partial H}{\partial p_j}, \quad j = 1, 2, \dots, n.$$

- A solution  $R^1(t) = (q_1^{(1)}(t), \dots, q_n^{(1)}(t), p_1^{(1)}(t), \dots, p_n^{(1)}(t))$  lives on a surface in  $2n$ -dimensional phase space.
- Chaotic system – exponential spreading:



$$|R^1(t) - R^2(t)| \propto e^{\lambda t}, \quad \lambda > 0.$$

Lyapunov exponent  $\lambda$

# Classical chaos

## KAM Theory

- Regular motion: Any solution  $R^1(t) = (q_1^{(1)}(t), \dots, q_n^{(1)}(t), p_1^{(1)}(t), \dots, p_n^{(1)}(t))$  lives on a tori in the  $2n$ -dimensional phase space.
- Add a perturbation  $V$  that breaks integrability. KAM describes how the tori is gradually deformed.
- Cranking up  $V$ : Going from regular to full blown chaos.



Poincaré section

# Quantum chaos

## Butterfly effect

*Schrödinger equation:*

$$\frac{d\hat{\rho}}{dt} = i[\hat{\rho}, \hat{H}].$$

- Trace distance

$$T(\hat{\rho}_1(t), \hat{\rho}_2(t)) \equiv \frac{1}{2} \text{Tr} \left[ \sqrt{(\hat{\rho}_1(t) - \hat{\rho}_2(t))^2} \right] = \frac{1}{2} \sum_i |\mu_i| \quad \stackrel{\text{Unitary linear evolution}}{=} \text{const.}$$

$\mu_i$  eigenvalues of  $(\hat{\rho}_1(t) - \hat{\rho}_2(t))$

- Quantum mechanics – linear theory.
- No Butterfly effect! Or...

# Quantum chaos

## Butterfly effect

- Perturbation  $\hat{\Gamma}$ :  $\hat{H}_1 = \hat{H}$  and  $\hat{H}_2 = \hat{H} + \hat{\Gamma}$ .
- Evolution,  $\frac{d\hat{\rho}_1}{dt} = i[\hat{\rho}_1, \hat{H}_1]$  and  $\frac{d\hat{\rho}_2}{dt} = i[\hat{\rho}_2, \hat{H}_2]$ .

- Trace distance

$$T(\hat{\rho}_1(t), \hat{\rho}_2(t)) \propto e^{\lambda t}$$


- Quantum butterfly effect!
- Non-unitary evolution  $\rightarrow$  butterfly effect,

$$\frac{d\hat{\rho}}{dt} = i[\hat{\rho}, \hat{H}] + \hat{L}[\hat{\rho}].$$

# Quantum chaos

## Characteristics of quantum chaotic systems

- Spectrum  $E_n$ .
- Energy separation  $s_n = E_{n+1} - E_n$ .
- Normalized distribution  $P(S)$ .
- Regular motion:  $P(S) = e^{-S}$  (Poisson distribution).
- Chaotic motion:  $P(S) = \frac{\pi}{2} S^\beta e^{-\pi S^2/4}$  (Wigner distribution).




Level repulsion



# Quantum chaos

## Characteristics of quantum chaotic systems

- Level repulsion  $\rightarrow$  varying time-scales.
- Level repulsion  $\rightarrow$  ergodicity.
- Level repulsion  $\rightarrow$  avoided crossings.



Driven Rabi model

# Chaos vs thermalization

## Spin-orbit coupled particle


Rabi Model

- Mean-field for the bosons, parametrize the atom by

$$|\theta\rangle = \begin{bmatrix} \sqrt{(1+Z)/2} \\ \sqrt{(1-Z)/2} e^{i\delta} \end{bmatrix},$$

- Semi-classical Hamiltonian

$$H_{cl} = \frac{p^2}{2} + \frac{x^2}{2} + \frac{\omega}{2} Z + (gx\sqrt{2} + \gamma)\sqrt{1-Z^2} \cos \delta.$$



Poincaré section.

- This Hamiltonian is chaotic in a classical sense  $\rightarrow$  thermalization.

# Chaos vs thermalization

Jahn-Teller Model

## Spin-orbit coupled particle

- 2D SO coupling.

$$\hat{H}_{SO} = \frac{\hat{p}^2}{2m} + \frac{1}{2}m\omega^2\hat{r}^2 + v_x\hat{p}_x\hat{\sigma}_x + v_y\hat{p}_y\hat{\sigma}_y.$$

- $\omega = 0 \rightarrow$  dispersions


$$E_{\pm}(p_x, p_y) = \frac{1}{2m}(p_x^2 + p_y^2) \pm \sqrt{(v_x p_x)^2 + (v_y p_y)^2}.$$

# Chaos vs thermalization


Jahn-Teller Model

## Spin-orbit coupled particle

- $v_x = v_y \rightarrow U(1)$  symmetry  
 $[\hat{J}, \hat{H}_{SO}] = 0$ .
- $v_x = v_y$  and  $\omega \neq 0 \rightarrow \hat{H}_{SO}$  equals dual  $E \times \varepsilon$  –Jahn-Teller model.



- $v_x \neq v_y \rightarrow Z_2$  symmetry  $[\hat{J}, \hat{H}_{SO}] \neq 0$ .
- $\hat{H}_{SO}$  equals dual  $E \times (\beta_1 + \beta_2)$  –Jahn-Teller model.





# Chaos vs thermalization

Jahn-Teller Model

Classical dynamics

- Poincaré sections ( $v_x \neq v_y$ ).



**Classical chaos**

# Chaos vs thermalization


Jahn-Teller Model

## Quantum dynamics

- Distributions ( $v_x \neq v_y$ ).

Quantum Scars  
Remnants of periodic classical solutions.

Heller, *Phys. Rev. Lett.* (1984).



Full Quantum

Truncated Wigner  
(Semi-classical)



**Ergodic and thermalization**

# Chaos vs thermalization

Jahn-Teller Model

## KAM theory

- "Islands" may survive large integrability breaking perturbations.




Poincaré sections

# Chaos vs thermalization

Jahn-Teller Model

## KAM theory

- Initiate a state in one island.



Distribution after long time when initial state in a regular island.

- No thermalization: Not all eigenstates obey *ETH*.





# Localization


# ETH revisited

## Ergodicity

- Thermalization  $\rightarrow$  ergodicity.
- Quantum information spreads over the whole accessible phase space.
- The information about a subsystem  $A$  is shared in the whole system  $S$ :

$\hat{\rho}_A(t)$  diagonal/mixed.


- $\hat{\rho}_A(t)$  obeys a "volume law".
- Can ergodicity be lost in quantum non-integrable/chaotic systems?



# Anderson localization

## Quantum interference

- Add disorder to your system.
- Time inversion symmetry.
- Enhance probability to scatter into the same state (factor 2) than an arbitrary state.
- Quantum interference effect. No counterpart in classical systems (particles).
- Strong localization, higher order interferences, vanishing conductivity.
- Non-ergodic, no thermalisation,  $\hat{\rho}_A(t)$  obeys area-law.



# Localization vs thermalization


- Spin models good for studying many-body localization

$$\hat{H} = \sum_i (J_x \hat{\sigma}_i^x \hat{\sigma}_{i+1}^x + J_y \hat{\sigma}_i^y \hat{\sigma}_{i+1}^y + J_z \hat{\sigma}_i^z \hat{\sigma}_{i+1}^z + h_i \hat{\sigma}_i^z)$$

- Clean *XXX* and *XXY* solvable, *XYZ* + *h* non-integrable.
- Localization with strong enough disorder  $h_i \in [-W, +W]$ .
- Localized eigenstates are not thermal, no thermalization!


Lieb-Robinson

Magnetization



# Many-body localization

- Does interaction destroy localization, i.e. giving thermalization? - No! (In general not, but sometimes)
- “Localization in Fock space”, Anderson problem on a hypercubic lattice.
- What are the loops causing localization?



**Hypercube:** Each vertex is connected to four other ones. MBL infinite number of edges!

**Thanks!**

